arXiv:1404.7400v1 [math.NT] 16 Feb 2014

THE EVALUATION OF THE SUMS OF MORE GENERAL SERIES BY BERNSTEIN POLYNOMIALS

Mehmet Acikgoz*, Ilknur Koca and Serkan Araci

University of Gaziantep, Faculty of Science and Arts, Department of Mathematics, 27310 Gaziantep, TURKEY

E-Mails: acikgoz@gantep.edu.tr; ibaltaci@gantep.edu.tr; mtsrkn@gmail.com
***Corresponding Author**

Abstract

Let n, k be the positive integers, and let $S_k(n)$ be the sums of the k-th power of positive integers up to n: $S_k(n) = \sum_{l=1}^{n} l^k$. By means of that we consider the evaluation of the sum of more general series by Bernstein polynomials. Additionally we show the reality of our idea with some examples.

2010 Mathematics Subject Classification. Primary 11B68, 11S80; Secondary 11M06.

Key Words and Phrases. Bernoulli numbers and polynomials, Bernstein polynomials, Sums of powers of integers.

1. Introduction

The history of Bernstein polynomials depends on Bernstein in 1904. It is well known that Bernstein polynomials play a crucial important role in the area of approximation theory and the other areas of mathematics, on which they have been studied by many researchers for a long time [1, 3, 5-7, 10, 11, 16, 17]. These polynomials also take an important role in physics.

Recently the works including applications of umbral calculus to Genocchi numbers and polynomials [2], the Legendre polynomials associated with Bernoulli, Euler, Hermite and Bernstein polynomials [3], the applications of umbral calculus to extended Kim's p-adic q-deformed fermionic integrals in the p-adic integer ring [4], the integral of the product of several Bernstein polynomials [5], the generating function of Bernstein polynomials [6], a theorem concerning Bernstein polynomials [10], new generating function of the (q-) Bernstein type polynomials and their interpolation function [11], q-analogues of the sums of powers of consecutive integers, squares, cubes, quarts and quints [12-15, 18-20], have been investigated extensively.

In the complex plane, the Bernoulli polynomials $B_n(x)$ are known by the following generating series:

$$\sum_{n=0}^{\infty} B_n(x)\frac{t^n}{n!} = \frac{t}{e^t-1}e^{xt},\ |t|<2\pi. \tag{1.1}$$

In the case $x=0$ in (1.1), we have $B_n(0) := B_n$ that stands for Bernoulli numbers. By (1.1), we have

$$B_n(x) = \sum_{k=0}^{n} \left(\begin{array}{c} n \\ k \end{array} \right) B_k x^{n-k}. \tag{1.2}$$

The Bernoulli numbers satisfy the following identity

$$B_0 = 1 \text{ and } (B+1)^n - B_n = \delta_{1,n}$$

where $\delta_{1,n}$ stands for Kronecker's delta and we have used $B^n := B_n$ (for details, see [3], [7], [9], [17]).

Recently, Acikgoz and Araci has constructed the generating function for the Bernstein polynomials $B_{k,n}(x)$ by the following rule:

$$\sum_{n=k}^{\infty} B_{k,n}(x)\frac{t^n}{n!} = \frac{(tx)^k}{k!}e^{t(1-x)}\ \ (t\in\mathbb{C} \text{ and } k=0,1,2,\cdots,n). \tag{1.3}$$

By (1.3), we see that

$$\sum_{n=k}^{\infty} B_{k,n}(x)\frac{t^n}{n!} = \sum_{n=k}^{\infty} \left(\left(\begin{array}{c} n \\ k \end{array} \right) x^k (1-x)^{n-k} \right) \frac{t^n}{n!}$$

by comparing the coefficients of $\frac{t^n}{n!}$ in the above, we derive well known expression of Bernstein polynomials, as follows: For $k,n\in Z_+$

$$B_{k,n}(x) = \left(\begin{array}{c} n \\ k \end{array} \right) x^k (1-x)^{n-k} \tag{1.4}$$

where $x\in[0,1]$ and $\left(\begin{array}{c} n \\ k \end{array} \right)$ is known as

$$\left(\begin{array}{c} n \\ k \end{array} \right) = \left\{ \begin{array}{cc} \frac{n!}{k!(n-k)!} & , \text{ if } n\geq k \\ 0 & , \text{ if } n<k. \end{array} \right. .$$

It follows from (1.4) that a few Bernstein polynomials are as follows:

$$B_{0,0}(x) = 1, B_{0,1}(x) = 1-x, B_{1,1}(x) = x, B_{0,2}(x) = (1-x)^2, B_{1,2}(x) = 2x(1-x)$$
$$B_{2,2}(x) = x^2, B_{0,3}(x) = (1-x)^3, B_{1,3}(x) = 3x(1-x)^2, B_{2,3}(x) = 3x^2(1-x), B_{3,3}(x) = x^3.$$

In the same time, the Bernstein polynomials $B_{k,n}(x)$ have several properties of interest:

- $B_{k,n}(x)\geq 0$, for $0\leq x\leq 1$ and $k=0,1,...,n$
- Bernstein polynomials have the symmetry property $B_{k,n}(x) = B_{n-k,n}(1-x)$
- $\sum_{k=0}^{n} B_{k,n}(x) = 1$, which is know a part of unity.
- $B_{k,n}(x) = (1-x)B_{k,n-1}(x) + xB_{k-1,n-1}(x)$ with $B_{k,n}(x) = 0$ for $k<0$, $k>n$ and $B_{0,0}(x) = 1$ *cf.* [1], [3], [5], [6], [7], [10], [16], [17].

From (1.1), a few Bernoulli polynomials can be generated as

$$B_0(x) = 1, B_1(x) = x - \frac{1}{2}, B_2(x) = x^2 - x + \frac{1}{6}, B_3(x) = x^3 - \frac{3}{2}x^2 + \frac{1}{2}x.$$

For any positive integer n, followings are the most known first three sums of powers of integers:

$$1 + 2 + 3 + ... + n = \frac{n(n+1)}{2},$$

$$1^2 + 2^2 + 3^2 + ... + n^2 = \frac{n(n+1)(2n+1)}{6}$$

and

$$1^3 + 2^3 + 3^3 + ... + n^3 = (1 + 2 + 3 + ... + n)^2 = \left[\frac{n(n+1)}{2}\right]^2.$$

Formulas for sums of integer powers were first given in generalizable form by mathematician Thomas Harriot (c. 1560-1621) of England. At about the same time, Johann Faulhaber (1580-1635) of Germany gave formulas for these sums, but he did not make clear how to generalize them. Also Pierre de Fermat (1601-1665) and Blaise Pascal (1623-1662) gave the formulas for sums of powers of integers.

The Swiss mathematician Jacob Bernoulli (1654-1705) is perhaps best and most deservedly known for presenting formulas for sums of integer powers. Because he gave the most explicit sufficient instructions for finding the coefficients of the formulas [12-15, 18-20].

So, we are interested in finding a method to derive a formula for the sums of powers of integers. Following an idea due to J. Bernoulli, we aim to obtain a Theorem which gives the method for the evaluation of the sums of more general series by Bernstein polynomials.

2. The Evaluation of the Sums of More General Series by Bernstein Polynomials

In the $17th$ century a topic of mathematical interest was finite sums of power of integers such as the series $1 + 2 + 3 + \cdots + (n-1)$ or the series $1^2 + 2^2 + 3^2 + \cdots + (n-1)^2$. The closed form for these finite sums were known, but the sums of the more general series $1^k + 2^k + 3^k + ... + (n-1)^k$ was not. It was the mathematician Jacob Bernoulli who would solve this problem with the following equality [12-15, 18-20]. The sum of the k-th powers of the first $(n-1)$ integers is given by the formula

$$1^k + 2^k + 3^k + ... + (n-1)^k = \int_1^n B_k(x)\, dx \tag{2.1}$$

using the integral of the Bernoulli polynomials, $B_n(x)$, under integral from 1 to n.

We are now in a position to express our aim as Theorem 1 for the evaluation of the sum of more general series by Bernstein polynomials.

Theorem 1. *Let n, k and m be positive integers and let $S_m(n)$ be $\sum_{l=1}^{n} l^m$, then we have*

$$S_m(n) = \frac{(-n^{-1})^k}{(m+k+1)!} \sum_{l=k}^{m+k+1} \binom{m+k+1}{l} B_{m+k-l+1} B_{k,l}(-n) - \frac{1}{(m+1)!} \sum_{l=0}^{m+1} \binom{m+1}{l} 2^{m+1-l} B_l + 1.$$

Proof. To prove this Theorem, we take $\sum_{k=0}^{\infty}\frac{t^k}{k!}$ in the both sides of the Eq. (2.1), so it yields to

$$\begin{aligned} e^t+e^{2t}+\cdots+e^{(n-1)t} &= \int_1^n\left(\sum_{k=0}^{\infty}B_k(x)\frac{t^k}{k!}\right)dx \\ &= \int_1^n\left[\frac{t}{e^t-1}e^{xt}\right]dx \\ &= \left[\sum_{m=0}^{\infty}B_m\frac{t^{m-1}}{m!}\right]\left[e^{nt}-e^t\right] \\ &= \left[\sum_{m=0}^{\infty}B_m\frac{t^{m-1}}{m!}\right]\left[\frac{n^{-k}(-1)^k k!}{e^t}\sum_{m=k}^{\infty}B_{k,m}(-n)\frac{t^{m-k}}{m!}-e^t\right] \end{aligned}$$

from the last identity, we see that

$$e^{2t}+e^{3t}+\cdots+e^{nt}=\frac{1}{t}\left[\sum_{m=0}^{\infty}B_m\frac{t^m}{m!}\right]\left[\frac{n^{-k}(-1)^k k!}{t^k}\sum_{m=0}^{\infty}B_{k,m}(-n)\frac{t^m}{m!}-\sum_{m=0}^{\infty}2^m\frac{t^m}{m!}\right] \quad (2.2)$$

by using Cauchy product rule in the right hand side of Eq. (2.2), we have

$$I_1=\sum_{m=0}^{\infty}\left(n^{-k}(-1)^k k!\sum_{l=k}^{m}\binom{m}{l}B_{m-l}B_{k,l}(-n)\right)\frac{t^{m-k-1}}{m!}-\sum_{m=0}^{\infty}\left(\sum_{l=0}^{m}\binom{m}{l}2^{m-l}B_l\right)\frac{t^{m-1}}{m!}.$$

By (2.2), we derive the following

$$I_2=\sum_{m=0}^{\infty}\left(2^m+3^m+\cdots+n^m\right)\frac{t^m}{m!}.$$

When we equate I_1 and I_2, we have

$$\begin{aligned} 1^m+2^m+3^m+\cdots+n^m &= \frac{\left(-n^{-1}\right)^k}{(m+k+1)!}\sum_{l=k}^{m+k+1}\binom{m+k+1}{l}B_{m+k-l+1}B_{k,l}(-n) \\ &\quad -\frac{1}{(m+1)!}\sum_{l=0}^{m+1}\binom{m+1}{l}2^{m+1-l}B_l+1. \end{aligned}$$

Thus, we complete the proof of the Theorem. □

Let $m=k$ in Theorem 1, we readily get the following Corollary 1.

Corollary 1. *Let n and k be positive integers and let $S_k(n)$ be $\sum_{l=1}^{n}l^k$, then we have*

$$S_k(n)=\frac{\left(-n^{-1}\right)^k}{(2k+1)!}\sum_{l=k}^{2k+1}\binom{2k+1}{l}B_{2k-l+1}B_{k,l}(-n)-\frac{1}{(k+1)!}\sum_{l=0}^{k+1}\binom{k+1}{l}2^{k+1-l}B_l+1.$$

Example 1. *Taking $k = 1$ in Corollary 1, we see that*

$$1 + 2 + 3 + ... + n = \frac{-n^{-1}}{6} \sum_{l=1}^{3} \binom{3}{l} B_{3-l} B_{1,l}(-n) - \frac{1}{2} \sum_{l=0}^{2} \binom{2}{l} 2^{2-l} B_l + 1$$
$$= \frac{n(n+1)}{2}.$$

For $k = 2$ in Corollary 1, we have

$$1^2 + 2^2 + 3^2 + ... + n^2 = \frac{n^{-2}}{120} \sum_{l=2}^{5} \binom{5}{l} B_{5-l} B_{2,l}(-n) - \frac{1}{6} \sum_{l=0}^{3} \binom{3}{l} 2^{3-l} B_l + 1$$
$$= \frac{n(n+1)(2n+1)}{6}.$$

By similar way, it can be easily shown for $k = 3, 4, \cdots$.

References

[1] S. Araci, *Novel identities for q-Genocchi numbers and polynomials,* Journal of Function Spaces and Applications, Volume **2012**, Article ID 214961, 13 pages, 2012.

[2] S. Araci, *Novel identities involving Genocchi numbers and polynomials arising from applications of umbral calculus*, Applied Mathematics and Computation (**2014**), http://dx.doi.org/10.1016/j.amc.2014.01.013 (In press).

[3] S. Araci, M. Acikgoz, A. Bagdasaryan, and E. Sen, *The Legendre polynomials associated with Bernoulli, Euler, Hermite and Bernstein polynomials*, Turkish Journal of Analysis and Number Theory, No. 1 (**2013**): 1-3. doi: 10.12691/tjant-1-1-1.

[4] S. Araci, M. Acikgoz, E. Sen, O*n the extended Kim's p-adic q-deformed fermionic integrals in the p-adic integer ring*, J. Number Theory 133 (**2013**), No.10, 3348-3361.

[5] M. Acikgoz And S. Araci, *A study on the integral of the Product of Several type Bernstein Polynomials*, IST Transactions of Applied Mathematics-Modeling and Simulation, Vol. 1, No. 1 (2) ISSN 1913-8342, pp. 10-14.

[6] M. Acikgoz and S. Araci, *On the generating function of the Bernstein polynomials,* Numerical Analysis and Applied Mathematics, **2010**, pp. 1141-1143.

[7] M. Acikgoz and S. Araci, *The relations between Bernoulli, Bernstein and Euler polynomials,* Numerical Analysis and Applied Mathematics, **2010**, pp. 1144-1147.

[8] M. Acikgoz and Y. Simsek, *On multiple interpolation functions of the Nörlund-type q-Euler polynomials*, Abstr. Appl. Anal. 2009, Art. ID 382574, 14 pages.

[9] G. S. Cheon, *A note on the Bernoulli and Euler polynomials*, Applied Mathematics Letters 16 (**2003**), 365-368.

[10] H. W. Gould, *A theorem concerning the Bernstein polynomials*, Math. Magazine 31 (5) (**1958**), 259-264.

[11] Y. Simsek and M. Acikgoz, *A new generating function of (q-) Bernstein type polynomials and their interpolation function*, Abstract and Applied Analysis, Volume 2010 (**2010**), Article ID 769095, 12 pages.

[12] Y. Simsek, D. Kim, T. Kim, S-H. Rim, *A note on the sums of powers of consecutive q-integers*, J. Appl. Funct. Differ. Equ. 1 (**2006**), No. 1, 81-88.

[13] Y. Simsek, T. Kim, S-H. Rim, *A note on the alternating sums of powers of consecutive q-integers*, Adv. Stud. Contemp. Math. 13 (**2006**), No. 2, 159-164.

[14] T. Kim, *Sums of powers of consecutive q-integers*, Adv. Stud. Contemp. Math. 9 (**2004**), 15-18.

[15] T. Kim, *A note on Exploring the sums of powers of consecutive q-integers*, Adv. Stud. Contemp. Math. 11 (2005), No. 1, pp. 137-140.

[16] T. Kim, *A note on q-Bernstein polynomials*, Russ. J. Math. Phys. 18(1), 73-82 (**2011**).
[17] M-S. Kim, T. Kim, B. Lee and C-S. Ryoo, *Some identities of Bernoulli numbers and polynomials associated with Bernstein polynomials*, Adv. Diff. Equa. Vol. **2010**, Article ID 305018, 7 pages.
[18] Y. -Y. Shen, *A note on the sums of powers of consecutive integers*, Tunghai Science 5 (**2003**), 101-106.
[19] M. Schlosser, *q-analogues of the sums of consecutive integers, squares, cubes, quarts and quints,* Electron. J. Combin. 11 (**2004**), #R71.
[20] K. C. Garrett and K. Hummel, *A combinatorial proof of the sum of q-cubes*, Electron. J. Combin.11 (**2004**).